Astronomy & Astrophysics manuscript no. Draft˙21.03.2013
March 22, 2013
© ESO 2013

arXiv:1303.5380v1 [astro-ph.HE] 21 Mar 2013

Rapid cooling of Cas A as a phase transition in dense QCD

Armen Sedrakian

Institute for Theoretical Physics, J. W. Goethe University, D-60438 Frankfurt am Main, Germany

March 22, 2013

ABSTRACT

We present a model of the compact star in Cassiopea A, which accommodates its unusually fast cooling behavior. This feature is interpreted as due to an enhancement in the neutrino emission triggered by a transition from fully gapped two-flavor color superconducting phase to a crystalline or an alternative gapless color superconducting phase. By fine-tuning a single parameter – the temperature of this transition - a specific cooling scenario can be selected that fits the Cas A data. Such scenario requires as a prerequisite a massive $M \sim 2M_\odot$ star and is, therefore, distinctive from models invoking canonical 1.4 $M_\odot$ mass star with nucleonic pairing only.

Key words. Neutron stars, Dense Matter, Pulsars

1. Introduction

Recently, an unprecedented fast cooling of the compact star in Cassiopeia A (Cas A) - the youngest known supernova remnant in Milky Way - was inferred from the Chandra observations over a period of 10 years (Heinke & Ho 2010; Shternin et al. 2011). The age of the object 330 yr is accurately deduced from the association of the remnant with the historical supernova SN 1680. This age estimate is confirmed through kinematical estimates. The fits to the surface temperature of the star suggest that the thermal soft x-ray spectrum originates from a non-magnetized carbon atmosphere at temperature $\sim 2. \times 10^6$ K and emitting region of 8-17 km.

Hadronic models of compact stars have been invoked to fit the data by assuming a canonical $1.4 M_\odot$ mass neutron star. The models proposed include those which attribute the Cas A cooling to a star that (*a*) undergoes a transition to superfluid state in the baryonic core, which is accompanied by an enhancement of the cooling due to Cooper pair-breaking processes (Page et al. 2011; Shternin et al. 2011); (*b*) experiences slower than commonly accepted thermal relaxation in the core due to low thermal conductivity (Blaschke et al. 2012); (*c*) undergoes rotationally induced compression which triggers a rapid Urca cooling in the center of the star (Negreiros et al. 2013).

Recent observations of PSR J1614-223010+11 lend firm evidence that equilibrium sequences of neutron stars contain massive $\sim 2M_\odot$ members (Demorest et al. 2010). Models which predict quark matter in such massive compact stars in color-superconducting state have been studied extensively (Alford et al. 2005a; Klähn et al. 2007; Ippolito et al. 2008; Knippel & Sedrakian 2009; Kurkela et al. 2010; Bonanno & Sedrakian 2012). Color superconductivity is important, because, otherwise, the unpaired quarks neutrino-cool a star rapidly to temperatures well below of those observed in Cas A (Page et al. 2000; Alford et al. 2005b; Anglani et al. 2006; Grigorian et al. 2005; Hess & Sedrakian 2011).

The true ground state of QCD at densities relevant for compact stars is not known. At relevant densities the strong coupling $\alpha_S \sim 1$, *i.e.*, perturbation theory is not useful. More importantly, because of β-equilibrium and strange quark mass, cross-flavor light, *i.e.*, up- and down-quark pairing occurs among the quarks living on different Fermi surfaces. As a consequence, there are modifications to the standard Bardeen-Cooper-Schrieffer (BCS) picture of pairing. The non-BCS pairing may lead to one of the possible gapless two-flavor phases (Shovkovy & Huang 2003; Müther & Sedrakian 2003; Shovkovy et al. 2005) or, for example, to the crystalline color-superconducting phase (Alford et al. 2001; Casalbuoni et al. 2005; Giannakis et al. 2005; Rajagopal & Sharma 2006; Gatto & Ruggieri 2007; Mannarelli et al. 2007; Sedrakian & Rischke 2009; Huang & Sedrakian 2010; Anglani et al. 2013). The crystalline-color-superconductivity comes in several variants, which differ by the way the translational symmetry is broken by the condensate of Cooper pairs carrying finite momentum. Below we shall use results obtained for the so-called Fulde-Ferrell (hereafter FF) phase, which is simple to model, but is general enough to preserve key feature of the crystalline phases, namely the existence of gapless modes on the Fermi surfaces of quarks (Alford et al. 2001; Sedrakian & Rischke 2009; Huang & Sedrakian 2010).

Proto-neutron stars emerge from supernova explosions with temperatures $T \sim 50$ MeV and nearly isospin symmetric matter in the core. After the initial fast cooling to $T \sim 0.1$ MeV the core reaches its asymptotic state which is highly isospin asymmetric. The initial fast temperature drop is followed by much slower cooling to temperatures of order 10 keV over the following $10^4 - 10^5$ years. Initially, high-temperature and low-isospin quark matter core is in the perfect two-flavor color-superconducting (hereafter 2SC) state where the Fermi surfaces of the up- and down-quarks are completely gapped. As the star cools and the β-equilibrium is established, the 2SC phase becomes unstable, because the asymptotic ground state of quark matter corresponds to a class of gapless and/or crystalline two-flavor color-superconducting phases mentioned above. This is illustrated in the top panel of Fig. 1, where we show the phase diagram of the FF and 2SC phases in the temperature and flavor asymmetry plane [adapted from Sedrakian & Rischke (2009)]. During the early evolution of the star, the matter traverses from the upper left corner of the phase diagram (i.e. from unpaired, almost isospin symmetric matter) to the right lower corner, which corresponds to the cold FF phase and large isospin asymmetry. The phase diagram of the same phases in the density-temperature



Fig. 1. Upper panel. The phase diagram of color-superconducting quark matter in the temperature (T) and isospin imbalance $\alpha = (n_d - n_u)/(n_u + n_d)$ plane, where n_d and n_u are the number densities of d and u flavors of quarks. The arrows show schematically the path in the phase diagram during the cooling of a star. Lower panel. Phase diagram of the same phases, but in the temperature and baryon density (n_B) plane for matter in β-equilibrium with electrons.

plane and under β-equilibrium is shown in the lower panel of Fig. 1 [adapted from Huang & Sedrakian (2010)].

In this work we conjecture that the rapid cooling of Cas A can be understood as a phase transition from the perfect 2SC phase to a crystalline/gapless color-superconducting state. Thus, somewhat counter-intuitively, the rapid cooling is the result of a phase transition *within* the phase diagram of dense QCD from one to another color-superconducting phase. This transition is characterized by a temperature T^* and the transient "widths" w, characterizing the finite time-scale of the transition.

In general T^* will depend on a number of factors, such as the nucleation history of the superconducting phase, geometry of the superconducting regions, dynamics and order of the phase transition, etc. Furthermore, because of the density dependence of the critical temperature T^* in the stellar interior, this phase transition will not occur globally in the quark core, but gradually within some shells. Therefore, the values of T^* and w will be determined by the local conditions in shells. We expect that the transition is not instantaneous: it takes place via a propagation of the phase-separation fronts rather than a coherent instantaneous transition. The *equilibrium* model calculations give $T_c \sim 10$ MeV in a particular (Nambu-Jona-Lasinio Lagrangian based) model, see Fig. 1. But the *absolute* value of the gap is strongly model dependent and it may differ from the equilibrium critical temperature of phase transition for dynamical reasons. (Supercooled state in electronic superconductor are observed, where the system remains in the initial unpaired state below the equilibrium critical transition temperature T_c, because of the finite time-scales of the phase transition process). These considerations motivate our treatment of T^* as a free parameter which will be used to fit our cooling model to the Cas A data. These fits are not sensitive to the precise value of transient width w characterizing the duration of the transition.

A preliminary report of the present study was given elsewhere (Sedrakian 2013).

2. Modeling cooling processes

Our cooling model is described in Hess & Sedrakian (2011), where the details of the microscopic input and the method of solution can be found. Here we describe the extension of this model, which allows us to accommodate the phase-transition between the phases in a phenomenological manner.

Quark matter consisting of two light u and d flavors of quarks cools predominantly via the beta decay (Urca) reactions $d \to u + e + \bar{\nu}$, $\quad u + e \to d + \nu$, where e stands for electron, ν and $\bar{\nu}$ represent electron neutrino and antineutrino. For unpaired quarks and to leading order in the strong coupling constant α_S the emissivity of the process *per quark color* was calculated by Iwamoto (1980)

$$\epsilon_\beta = \frac{914}{945} G^2 \cos^2\theta \, p_d p_u p_e \alpha_s T^6, \tag{1}$$

where G is the weak coupling constant, θ is the Cabibbo angle, p_d, p_u, and p_e are the Fermi momenta of down quarks, up quarks, and electrons.

Quark pairing modifies the temperature dependence of the process (1). In the BCS-type superconductors, the process is suppressed almost linearly for $T \simeq T_c$ and exponentially for $T \ll T_c$, where T_c is the critical temperature. In gapless superconductors the emergence of the new scale $\delta\mu = (\mu_d - \mu_u)/2$, where $\mu_{u,d}$ are the chemical potentials of light quarks, leads to two qualitatively different possibilities, which are distinguished by the value of the dimensionless parameter $\zeta = \Delta_0/\delta\mu$, where Δ_0 is the gap in the limit $\delta\mu = 0$ (Jaikumar et al. 2006). When $\zeta > 1$, the entire Fermi surface is gapped. Consequently, when the thermal smearing of the Fermi surface $\sim T$ is much smaller than the gap, no excitation can be created out of the Fermi sphere. When $\zeta < 1$, particles can be excited from the gapless regions of the Fermi sphere. For the FF phases the shift in the chemical potential is replaced by the anti-symmetric in flavor part of the single particle spectrum of up- and down-quarks, $\epsilon_{u/d}$, *i.e.*, $\delta\mu \to [\epsilon_d(\boldsymbol{Q}) - \epsilon_u(\boldsymbol{Q})]/2$, where $\boldsymbol{Q}$ is the total momentum of a Cooper pair. In the case of single plane wave crystalline phase the spectrum depends on the magnitude of the vector $\boldsymbol{Q}$ only. The asymptotic behavior of the ζ parameter discussed above, permits us to use an interpolation formula, which covers the asymptotic limits, and does not require detailed modeling of the



Fig. 2. Left panel. Dependence of the surface temperature on time for the values of the transition temperature T^* (in MeV) indicated in the plot and for fixed width $w = 0.2$ MeV. The underlying model is the 1.91 $M_\odot$ compact star model of Knippel & Sedrakian (2009). Right panel. Dependence of the surface temperature (in units of 10^6 K) on time for the same model. The points with error bars show the Cas A data, the solid lines are fits to these data by variation of T^* (again for fixed width $w = 0.2$ MeV).

physics of the underlying phase (Jaikumar et al. 2006)

$$\epsilon_\beta^{rg}(\zeta; T \le T_c) = 2f(\zeta)\epsilon_\beta, \quad f(\zeta) = \frac{1}{\exp[(\zeta - 1)\frac{\delta\mu}{T} - 1] + 1}, \quad (2)$$

where the factor two is the number of colors.

Because the 2SC pairing pattern breaks the SU(3) color symmetry, one of the quark colors (say, the blue color) is not involved in that pairing. The strength and the flavor content of pairing among blue quarks is model dependent (Alford et al. 2003; Schmitt et al. 2003; Schmitt 2005; Aguilera et al. 2005, 2006; Aguilera 2007). It is likely that cross-flavor (blue-up and blue-down) pairing is suppressed by large mismatch between the Fermi surfaces of the up and down quarks. Pairing is expected in the color $\mathbf{6}_S$ and flavor $\mathbf{3}_S$ channel, which is same-flavor and same-color pairing, and therefore is not affected by the flavor asymmetry (Alford et al. 2003). However, the pairing in this channel vanishes in the chiral limit, when the mass of blue quarks vanishes. Pairing due to alternative mechanisms, outside those studied so far is not excluded. Quite generally, single flavor and/or color pairings have gaps in the range of 10-100 keV, which are much smaller than the gaps emerging from 2SC-type pairing. Since there are no definitive theoretical prediction about the pairing in the blue sector we will assume that the neutrino emissivity of blue quarks is suppressed exponentially by the pairing gap, as is the case for nucleonic matter. Therefore, we shall simply parameterize the modifications of the Urca process on blue quarks as in the case of baryonic matter, *i.e.*,

$$\epsilon_\beta^{b}(T \le T_c) = \epsilon_\beta^{b} \exp\left(-\frac{\Delta_b}{T}\right), \quad (3)$$

where Δ_b is the pairing gap for the blue color quarks. Note that $\epsilon_\beta^b = \epsilon_\beta^{rg}/2$ due to the different number of colors involved. Several studies used similar parameterizations (Grigorian et al. 2005; Hess & Sedrakian 2011). Furthermore, we will choose to work with a pairing gap $\Delta_b = 100$ keV, which has the consequence that the blue quarks do not affect the cooling dynamics and the fits to Cas A data, *i.e.*, they play only a passive role. The effect of smaller values of this gap on the cooling is discussed elsewhere (Hess & Sedrakian 2011).

In addition to neutrino emissivities we need to model the specific heat of the quark phases. For the inhomogeneous red-green condensate the critical temperature changes with $\delta\mu$ (Hess & Sedrakian 2011)

$$T_c(\zeta) \simeq T_{c0}\sqrt{1 - \frac{4\mu}{3\Delta_0}\delta(\zeta)}, \quad (4)$$

where $\mu = (\mu_d + \mu_u)/2$, $\Delta_0 = \Delta(\zeta = 0)$, $T_{c0} = T_c(\zeta = 0)$, and $\delta(\zeta) = (n_d - n_u)/(n_d + n_u)$. The fully gapped and gapless regimes behave differently; in the presence of gapless modes, the specific heat has a linear dependence on the temperature. A phenomenological way to model this behavior is given by the relation

$$c_S^{rg}(\zeta; T \le T_c) = f(\zeta)c_N^{rg}, \quad (5)$$

where c_N^{rg} is the specific heat of red-green unpaired quarks, taken as that for noninteracting quarks and c_S^{rg} is the specific heat of pair-correlated quarks. To compute the specific heat of the electrons, we assume a noninteracting, ultrarelativistic gas.

Finally, we need the relation between the temperature of the isothermal core and the crust, which can be written as (Potekhin et al. 1997)

$$T_{s6} = (\alpha T_8)^\beta g_s^{1/4} \quad (6)$$

where T_{s6} is the surface temperature in units of 10^6 K, and T_8 is the isothermal core temperature in units of 10^8 K, α and β are constants which depend on the composition of the star's atmosphere, g_s is the surface gravity in units of 10^{14} cm s^{-2}. We will adopt the values $\beta = 0.55$, which is intermediate between the purely-iron ($\beta = 0.5$) and fully accreted envelope ($\beta = 0.61$) and $\alpha = 181$ appropriate for an accreted envelop. The fit results are sensitive to the first parameter, but not the second.

3. Simulation results

The key idea of our model is that the transition occurs from the fully gap (perfect) 2SC phase to some type of crystalline/gapless phase. A particular realization of this scenario would be a transition to single plane wave crystalline phase (the FF phase). We model this transition phenomenologically by adopting temperature dependent (and, therefore, time-dependent) parameter $\zeta(T)$. [Note that Hess & Sedrakian (2011) assumed constant ζ and thus phase transitions of the type discussed here were excluded in their study from the outset]. The temperature/time dependence is modelled such that for $T > T^*$ one obtains $\zeta > 1$, whereas for $T < T^*$, $\zeta < 1$. The $\zeta(T)$ function is modelled by the simply law

$$\zeta(T) = \zeta_i - \Delta\zeta \, g(T), \tag{7}$$

where ζ_i is the initial value, $\Delta\zeta$ is the constant change in this function, and the function $g(T)$ affects the transition from the initial value ζ_i to the asymptotic final value $\zeta_f = \zeta_i - \delta\zeta$. The transition is conveniently modelled by the following function

$$g(T) = \frac{1}{\exp\left(\frac{T-T^*}{w}\right) + 1}, \tag{8}$$

which allows us to control the temperature of transition, fixed by T^*, and the smoothness of the transition, fixed by the width w.

Fig. 2, left panel, shows the cooling evolution of $M = 1.91 M_\odot$ model of a compact star for several values of T^* and fixed $w = 0.2$ MeV. It is seen that for a specific value of T^* the cooling curve passes through the location of Cas A in the log T-log t plane. The right panel of Fig. 2 shows the observed evolution of Cas A for the same value of w and two (fine-tuned) values of T^*; the best fit to data is shown by solid line; the dashed line demonstrates the sensitivity of the fit to variations of the parameter T^*. In each case the critical temperature of the same-color pairing of the blue quarks satisfies the condition $T_{cb} = \Delta_b/1.76 > T^*$, which implies that blue quarks have no effect on the dynamics of the transition process. The numerical value $\Delta_b = 0.1$ MeV was used in the simulations. In case of low critical temperatures of blue quark pairing a compact star will rapidly neutrino-cool on time-scales $t \leq 300$ yr below the temperature of Cas A, which obviously will preclude any fit to the data.

4. Conclusions

We have demonstrated that the rapid cooling of Cas A can be understood as a phase transition from the perfect two-flavor color-superconducting phase to a gapless/crystalline phase. Counterintuitively, *the phase transition occurs within the phase diagram of dense QCD and reflects the ordering of various superconducting phases in the temperature, density and isospin spaces.* This idea is supported qualitatively by the structure of the phase diagram of the dense QCD at densities and temperatures relevant to cooling evolution of compact stars.

We have shown that the data can be fitted by adjusting a single parameter - the phase transition temperature - at a fixed value of transient time-scale, which encodes the dynamics of the transition. A further important assumption is that there is no interference from the blue quarks, so that the temperature of the model is above the Cas A temperature for $t \leq 300$ yr, which is an obvious requirement for any successful fit to data in the absence of internal heating.

The present model implies that only massive compact stars undergo a rapid temperature drop observed in Cas A, for only massive members of compact star sequences contain (color-superconducting) quark matter in their interiors. This feature sets our model apart from the alternatives which invoke nucleonic pairing and pair-breaking processes (Page et al. 2011; Shternin et al. 2011) or slow thermal relaxation (Blaschke et al. 2012), *i.e.*, physics of canonical mass neutron stars. Rotation is unimportant for our modeling as opposed to the model of Negreiros et al. (2013). Clearly, the various models can be discriminated through their distinctive features in case further observational information will become available on the properties of Cas A. Evidently, it would be premature to conclude that the Cas A behavior is a *direct evidence for superfluidity* or even *measurement of T_c for any particular type of a superfluid* in neutron star interiors.

Acknowledgments

The author is grateful to M. Alford, H. Grigorian, D. Hess, X.-G. Huang, K. Rajagopal, D. H. Rischke, A. Schmitt, and I. Wasserman for useful discussions and correspondance. This work is partially supported though a collaborative research grant of the Volkswagen Foundation (Hannover, Germany).

References

Aguilera, D. N. 2007, Ap&SS, 308, 443
Aguilera, D. N., Blaschke, D., Buballa, M., & Yudichev, V. L. 2005, Phys. Rev. D, 72, 034008
Aguilera, D. N., Blaschke, D., Grigorian, H., & Scoccola, N. N. 2006, Phys. Rev. D, 74, 114005
Alford, M., Bowers, J. A., & Rajagopal, K. 2001, Phys. Rev. D, 63, 074016
Alford, M., Braby, M., Paris, M., & Reddy, S. 2005a, ApJ, 629, 969
Alford, M., Jotwani, P., Kouvaris, C., Kundu, J., & Rajagopal, K. 2005b, Phys. Rev. D, 71, 114011
Alford, M. G., Bowers, J. A., Cheyne, J. M., & Cowan, G. A. 2003, Phys. Rev. D, 67, 054018
Anglani, R., Casalbuoni, R., Ciminale, M., et al. 2013, ArXiv e-prints
Anglani, R., Nardulli, G., Ruggieri, M., & Mannarelli, M. 2006, Phys. Rev. D, 74, 074005
Blaschke, D., Grigorian, H., Voskresensky, D. N., & Weber, F. 2012, Phys. Rev. C, 85, 022802
Bonanno, L. & Sedrakian, A. 2012, A&A, 539, A16
Casalbuoni, R., Gatto, R., Ippolito, N., Nardulli, G., & Ruggieri, M. 2005, Physics Letters B, 627, 89
Demorest, P. B., Pennucci, T., Ransom, S. M., Roberts, M. S. E., & Hessels, J. W. T. 2010, Nature, 467, 1081
Gatto, R. & Ruggieri, M. 2007, Phys. Rev. D, 75, 114004
Giannakis, I., Hou, D.-F., & Ren, H.-C. 2005, Physics Letters B, 631, 16
Grigorian, H., Blaschke, D., & Voskresensky, D. 2005, Phys. Rev. C, 71, 045801
Heinke, C. O. & Ho, W. C. G. 2010, ApJ, 719, L167
Hess, D. & Sedrakian, A. 2011, Phys. Rev. D, 84, 063015
Huang, X.-G. & Sedrakian, A. 2010, Phys. Rev. D, 82, 045029
Ippolito, N. D., Ruggieri, M., Rischke, D. H., Sedrakian, A., & Weber, F. 2008, Phys. Rev. D, 77, 023004
Iwamoto, N. 1980, Physical Review Letters, 44, 1637
Jaikumar, P., Roberts, C. D., & Sedrakian, A. 2006, Phys. Rev. C, 73, 042801
Klähn, T., Blaschke, D., Sandin, F., et al. 2007, Physics Letters B, 654, 170
Knippel, B. & Sedrakian, A. 2009, Phys. Rev. D, 79, 083007
Kurkela, A., Romatschke, P., Vuorinen, A., & Wu, B. 2010, ArXiv e-prints
Mannarelli, M., Rajagopal, K., & Sharma, R. 2007, Phys. Rev. D, 76, 074026
Müther, H. & Sedrakian, A. 2003, Phys. Rev. D, 67, 085024
Negreiros, R., Schramm, S., & Weber, F. 2013, Physics Letters B, 718, 1176

Page, D., Prakash, M., Lattimer, J. M., & Steiner, A. W. 2000, Physical Review Letters, 85, 2048
Page, D., Prakash, M., Lattimer, J. M., & Steiner, A. W. 2011, Physical Review Letters, 106, 081101
Potekhin, A. Y., Chabrier, G., & Yakovlev, D. G. 1997, A&A, 323, 415
Rajagopal, K. & Sharma, R. 2006, Phys. Rev. D, 74, 094019
Schmitt, A. 2005, Phys. Rev. D, 71, 054016
Schmitt, A., Wang, Q., & Rischke, D. H. 2003, Physical Review Letters, 91, 242301
Sedrakian, A. 2013, ArXiv e-prints
Sedrakian, A. & Rischke, D. H. 2009, Phys. Rev. D, 80, 074022
Shovkovy, I. & Huang, M. 2003, Physics Letters B, 564, 205
Shovkovy, I. A., Rüster, S. B., & Rischke, D. H. 2005, Journal of Physics G Nuclear Physics, 31, 849
Shternin, P. S., Yakovlev, D. G., Heinke, C. O., Ho, W. C. G., & Patnaude, D. J. 2011, MNRAS, 412, L108